

## POLYNOMIALS (10/11/17)

### EASIER PROBLEMS

1. The polynomial  $f(x) = x^n + a_1x^{n-1} + \cdots + a_n$  has integer coefficients and  $n$  distinct integer roots. Suppose that all of its roots are coprime. Show that  $a_n$  and  $a_{n-1}$  are coprime.
2.  $a$ ,  $b$ , and  $c$  are the three roots of the polynomial  $x^3 - 3x^2 + 1$ . Find  $a^3 + b^3 + c^3$ .
3.  $p(x)$  is a polynomial with integer coefficients such that  $p(0) = 1$ ,  $p(1) = 2$ ,  $p(-1) = 1$ . Prove that  $p(x)$  has no integer roots.
4. (AIME 1989) Assume that  $x_1, x_2, \dots, x_7$  are real numbers such that

$$\begin{aligned}x_1 + 4x_2 + 9x_3 + 16x_4 + 25x_5 + 36x_6 + 49x_7 &= 1 \\4x_1 + 9x_2 + 16x_3 + 25x_4 + 36x_5 + 49x_6 + 64x_7 &= 12 \\9x_1 + 16x_2 + 25x_3 + 36x_4 + 49x_5 + 64x_6 + 81x_7 &= 123.\end{aligned}$$

Find the value of

$$16x_1 + 25x_2 + 36x_3 + 49x_4 + 64x_5 + 81x_6 + 100x_7.$$

### PUTNAM PROBLEMS (NOT NECESSARILY HARDER)

5. (Putnam 2007) Let  $f$  be a polynomial with positive integer coefficients. Prove that if  $n$  is a positive integer, then  $f(n)$  divides  $f(f(n) + 1)$  if and only if  $n = 1$ .
6. (Putnam 2009) Let  $p(x)$  be a real polynomial that is nonnegative for all real  $x$ . Prove that for some  $k$ , there are real polynomials  $f_1(x), \dots, f_k(x)$  such that  $p(x) = \sum_{i=1}^k f_i(x)^2$ .
7. (Putnam 2003) Do there exist polynomials  $a(x), b(x), c(y), d(y)$  such that

$$1 + xy + x^2y^2 = a(x)c(y) + b(x)d(y)?$$

8. (Putnam 2010) Find all polynomials  $P(x), Q(x)$  with real coefficients such that  $P(x)Q(x+1) - P(x+1)Q(x) = 1$ .
9. (Putnam 2008) Let  $n \geq 3$  be an integer. Let  $f(x)$  and  $g(x)$  be polynomials with real coefficients such that the points

$$(f(1), g(1)), (f(2), g(2)), \dots, (f(n), g(n)) \in \mathbb{R}^2$$

are the vertices of a regular  $n$ -gon in counterclockwise order. Prove that at least one of  $f(x)$  and  $g(x)$  has degree greater or equal to  $n - 1$ . (By the way, suppose  $n$  is odd and  $\deg(f(x)) < n - 1$ . Can you say anything special about the regular  $n$ -gon?)

10. (Putnam 2005) Let  $p(z)$  be a polynomial of degree  $n$  all of whose zeros have absolute value 1 in the complex plane. Put  $g(z) = p(z)/z^{n/2}$ . Show that all zeros of  $g'(z)$  have absolute value 1.

## GOOD THINGS TO KNOW

- The Fundamental Theorem of Algebra.
- Expressions for coefficients in terms of roots (as elementary symmetric polynomials).
- Roots of real polynomials (must occur in conjugate pairs).
- Intermediate Value Theorem.
- Mean Value Theorem: For a real-valued function, the derivative must vanish between two zeros of the function.
- Divisibility: for an integer polynomial  $p(x)$ ,  $(a - b) \mid (p(a) - p(b))$ .

## HINTS

1. Think about how  $a_i$ 's are expressed in terms of roots.
2. Express  $a^3 + b^3 + c^3$  in terms of the elementary symmetric polynomials.
3. What is  $p(x)$  modulo 3?
4. The left-hand sides are values of a (low-degree!) polynomial.
5. Not much to say - straight divisibility question.
6. What does the factorization of  $p(x)$  look like?
7. Consider the coefficients of  $y^k$  on the right-hand side as polynomials of  $x$ . How are they related to  $a(x)$  and  $b(x)$ ? (Or plug in particular values of  $y$  into both sides.)
8. Compare  $P(x)(Q(x + 1) - Q(x))$  and  $Q(x)(P(x + 1) - P(x))$ .
9. What can you say about the vector-valued polynomial function  
 $(f(x + 1) - f(x), g(x + 1) - g(x))$   
?  
10. What is special about the values of  $g(z)$  if  $|z| = 1$ ? Also, think about the Mean Value Theorem.