

Binomial coefficients and generating functions—Answers and Hints

February 16, 2020

Binomial coefficients

The binomial coefficients $\binom{n}{k}$ counts the number of ways one can choose k objects from given n . They are coefficients in the binomial expansion

$$(x+1)^n = \binom{n}{0}x^n + \binom{n}{1}x^{n-1} + \cdots + \binom{n}{n-1}x + \binom{n}{n}.$$

More explicitly,

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} = \frac{n(n-1)\cdots(n-k+1)}{k!}.$$

A very useful formula for the binomial coefficients is

$$\binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1}. \quad (1)$$

Here are two simple exercises about binomial coefficients.

- Prove that if $n = 2^m$ with m a positive integer, then

$$\binom{n}{k}$$

is an even integer for any $1 \leq k \leq n-1$.

Hint: Use the formula

$$\binom{n}{k} = n/k \binom{n-1}{k-1}. \quad (2)$$

- Let m and n be integers such that $1 \leq m \leq n$. Prove that m divides the number

$$n \sum_{k=0}^{m-1} (-1)^k \binom{n}{k}.$$

Hint: Use formula (1) to simplify the sum, then use formula (2).

Generating functions

The terms of a sequence $(a_n)_{n \geq 0}$ can be combined into a function

$$G(x) = a_0 + a_1x + a_2x^2 + \cdots + a_nx^n + \cdots,$$

called the generating function of the sequence. For example, the finite sequence $\binom{m}{n}$, with m fixed and n varies, gives the function $(x+1)^m$. The generating function for $a_n = \frac{1}{n}$ is $-\ln(1-x)$.

Generating functions provide a method to understand recursive relations of a sequence.

Theorem. Suppose a_n ($n \geq 0$) is a sequence satisfying a second-order linear recurrence,

$$a_n + ua_{n-1} + va_{n-2} = 0.$$

Suppose that the quadratic equation $\lambda^2 + u\lambda + v = 0$ has two distinct roots r_1, r_2 . Then

$$a_n = \alpha r_1^n + \beta r_2^n$$

for some real numbers r_1, r_2 .

Proof. Let $G(x) = a_0 + a_1x + a_2x^2 + \dots$ be the generating function of a_n . The recurrence relation of a_n implies that

$$G(x) - a_0 - a_1x + u(G(x) - a_0) + vx^2G(x) = 0.$$

Solving for $G(x)$, we have

$$G(x) = \frac{a_0 + (ua_0 + a_1)x}{1 + ux + vx^2} = \frac{a_0 + (ua_0 + a_1)x}{(1 - r_1x)(1 - r_2x)}.$$

Using partial fractions, we have

$$G(x) = \frac{a_0 + (ua_0 + a_1)x}{(1 - r_1x)(1 - r_2x)} = \frac{\alpha}{1 - r_1x} + \frac{\beta}{1 - r_2x} = \sum_{n=1}^{\infty} (\alpha r_1^n + \beta r_2^n)x^n.$$

Therefore, $a_n = \alpha r_1^n + \beta r_2^n$. □

1. Prove the identity

$$\binom{m+n}{k} = \sum_{j=0}^k \binom{m}{j} \binom{n}{k-j}.$$

Hint: $(x+1)^{m+n} = (x+1)^m(x+1)^n$.

2. Give two proofs of the identity

$$\sum_{j=0}^n 2^{n-j} \binom{n}{j} \binom{j}{\lfloor j/2 \rfloor} = \binom{2n+1}{n}.$$

Hint: $(x+1)^{2n+1} = ((x+1)^2)^n(x+1)$.

Problems

1. Prove the identity

$$\sum_{k=1}^n k \binom{n}{k}^2 = n \binom{2n-1}{n-1}.$$

Putnam and beyond 880.

2. Find the general formula for the sequence $(y_n)_{n \geq 0}$ with $y_0 = 1$ and $y_n = ay_{n-1} + b^n$ for $n \geq 1$, where a and b are two fixed distinct real numbers. **Putnam and beyond 867.**
3. Prove that the Fibonacci numbers F_n satisfy

$$F_n = \binom{n}{0} + \binom{n-1}{1} + \binom{n-2}{2} + \dots.$$

Hint: use induction. **Putnam and beyond 872.**

4. Consider the triangular $n \times n$ matrix

$$A = \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 \\ 0 & 1 & 1 & \cdots & 1 \\ 0 & 0 & 1 & \cdots & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}.$$

Compute the matrix A^k for $k \geq 1$.

Putnam and beyond 860.

5. Show that the coefficient of x^k in the expansion of $(1 + x + x^2 + x^3)^n$ is

$$\sum_{j=0}^k \binom{n}{k} \binom{n}{k-2j}.$$

Putnam 1992 B-2

6. Find

$$\binom{n}{1}1^2 + \binom{n}{2}2^2 + \binom{n}{3}3^2 + \cdots + \binom{n}{n}n^2.$$

Hint: use formula (2) twice. **Putnam 1962 A-5**

7. Let S_0 be a finite set of positive integers. We define finite sets S_1, S_2, \dots of positive integers as follows. The integer a is in S_{n+1} if and only if exactly one of $a-1$ or a is in S_n . Show that there are infinitely many integers N for which

$$S_N = S_0 \cup \{N + a \mid a \in S_0\}.$$

Putnam 2000 B-5

8. For a set S of nonnegative integers, let $r_S(n)$ denote the number of ordered pairs (s_1, s_2) such that

$$s_1, s_2 \in S, s_1 \neq s_2, \quad \text{and} \quad s_1 + s_2 = n.$$

Is it possible to partition the nonnegative integers into two sets A and B in such a way that $r_A(n) = r_B(n)$ for all n ?

Putnam 2003 A-6

9. For positive integer n , denote by $S(n)$ the number of choices of the signs “+” or “−” such that $\pm 1 \pm 2 \pm \cdots \pm n = 0$. Prove that

$$S(n) = \frac{2^{n-1}}{\pi} \int_0^{2\pi} \cos t \cos 2t \cdots \cos nt \, dt.$$

Hint: write $\cos kt$ as $(e^{\sqrt{-1}kt} + e^{-\sqrt{-1}kt})/2$.

10. Let S_n denote the set of all permutations of the numbers $1, 2, \dots, n$. For $\pi \in S_n$, let $\sigma(\pi) = 1$ if π is an even permutation and $\sigma(\pi) = -1$ if π is an odd permutation. Also, let $\nu(\pi)$ denote the number of fixed points of π . Show that

$$\sum_{\pi \in S_n} \frac{\sigma(\pi)}{\nu(\pi) + 1} = (-1)^{n+1} \frac{n}{n+1}.$$

Putnam 2005 B-6